

Important information for the patient VANFLYTA can cause an abnormal electrical activity in your heart called "prolonged QT interval" which may lead to a life-threatening disturbance of the heart rhythm. Your healthcare professional will check the electrical activity of your heart with a test called an electrocardiogram (ECG) before you take VANFLYTA and while you are taking it. Contact your doctor immediately if: - You are feeling dizzy, lightheaded, or faint. - You sense a change in heart rhythm, for example,	palpitations or an abnormality of your pulse. You may feel your heart is beating too fast, but you may also sense a more nonspecific or vague change. - You have fainted or have been unconscious, even if it was only for a very short period of time, for example, seconds. - You suffer from diarrhea or vomiting, or are unable to eat or drink fluids in sufficient amounts. - You feel any other sudden change in your well-being. - Your medicines are changed by a physician other than the prescribing physician for VANFLYTA.	Consult your doctor first before taking VANFLYTA with any other medicines, including medicines obtained without a prescription or supplements as these may increase the risk of you developing QT interval prolongation. For more information, please read the package leaflet. Find the full product monograph that is prepared for healthcare professionals and includes Patient Medication Information by visiting the Health Canada website: https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html; the manufacturer's website www.daiichi-sankyo.ca, or by calling 1-866-791-9292.	Reporting Side Effects You can report any suspected side effects associated with the use of health products to Health Canada by: - Visiting the Web page on Adverse Reaction Reporting (canada.ca/drug-device-reporting) for information on how to report online, by mail or by fax; or - Calling toll-free at 1-866-234-2345. <i>NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.</i>  Daiichi-Sankyo	PATIENT CARD VANFLYTA® quizartinib - Please keep this card with you at all times. - This card contains important safety information that you should know before you take VANFLYTA and during treatment with VANFLYTA. - Show this card to any doctor, nurse, pharmacist or surgeon before any medical intervention or treatment.
			CA 122977 	 Please detach this section here.

Patient information Patient name: Date of Birth: In case of emergency, please contact: Name: Phone number:	Treatment information (To be completed by physician or patient) VANFLYTA has been prescribed at a once-daily dose of: Started on: mg / (mm/yy)	Prescriber information (To be completed by physician or patient) For more information or in case of emergency, please contact: Physician's name: Phone number:	Important information for healthcare professionals VANFLYTA can prolong the QT interval. Torsades de pointes, cardiac arrest, and sudden death have been reported in patients receiving VANFLYTA. - Reduce VANFLYTA dose if QTcF is >480 ms, interrupt VANFLYTA if QTcF is >500 ms, and permanently discontinue if associated with torsades de pointes, polymorphic ventricular tachycardia or signs/symptoms of life-threatening arrhythmia. - During VANFLYTA treatment, monitor ECG and serum electrolytes;	correct hypokalemia and hypomagnesemia. - Avoid non-essential medicines that prolong the QT interval. If unavoidable, monitor ECG frequently. - The dose of VANFLYTA should be reduced when used concomitantly with strong CYP3A4/5 inhibitors. For more information, please see the Product Monograph. VANFLYTA® is a registered trademark of Daiichi Sankyo Co., Ltd., used by Daiichi Sankyo Pharma Canada Ltd. under license. Copyright © 2025, Daiichi Sankyo Pharma Canada Ltd.